

RAPIDLY ROTATING STRANGE STARS FOR A NEW EQUATION OF STATE OF STRANGE QUARK MATTER

IGNAZIO BOMBACI¹, ARUN V. THAMPAN², AND BHASKAR DATTA^{3,4}*Draft version February 1, 2008*

ABSTRACT

For a new equation of state of strange quark matter, we construct equilibrium sequences of rapidly rotating strange stars in general relativity. The sequences are the normal and supramassive evolutionary sequences of constant rest mass. We also calculate equilibrium sequences for a constant value of Ω corresponding to the most rapidly rotating pulsar PSR 1937 +21. In addition to this, we calculate the radius of the marginally stable orbit and its dependence on Ω , relevant for modeling of kilo-Hertz quasi-periodic oscillations in X-ray binaries.

Subject headings: stars: neutron - equation of state - general relativity

1. INTRODUCTION

One of the most fascinating aspects of modern astrophysics, is the possible existence of a new family of collapsed stars consisting completely of a deconfined mixture of *up* (*u*), *down* (*d*), and *strange* (*s*) quarks, together with an appropriate number of electrons to guarantee electrical neutrality. Such compact stars have been referred to in the literature, as strange stars (SS), and their constituent matter as strange quark matter (SQM). The possible existence of SS is a direct consequence of the the so called *strange matter hypothesis*, formulated by Witten (1984) (see also Bodmer 1971, Terazawa 1979). According to this hypothesis, SQM, in equilibrium with respect to the weak interactions, could be the absolute ground state of strongly interacting matter rather than ⁵⁶Fe. Ever since the formulation of this hypothesis, there have been several reports in the literature, on the structure of non-rotating SS (e.g. Haensel, Zdunik & Schaefer 1986; Alcock, Farhi & Olinto 1986). However, these have remained largely theoretical and speculative in nature. Observational data from RXTE of certain low mass X-ray binaries (LMXBs) have recently given rise to a resurgence in the astrophysical interest associated with SS. It has been suggested that at least a few LMXBs could be harboring SS. For example, recent studies have shown that the compact objects associated with the X-ray burster 4U 1820 –30 (Bombaci 1997), the bursting X-ray pulsar GRO J1744 –28 (Cheng *et al.* 1998) and the X-ray pulsar Her X–1 (Dey *et al.* 1998) are likely SS candidates. The most promising and convincing SS candidates are the compact objects in the bursting millisecond X-ray pulsar SAX J1808.4 –3658 (Li *et al.* 1999a), and in the atoll source 4U 1728 –34 (Li *et al.* 1999b).

The compact nature of these sources make general relativity important in describing these systems. Furthermore, their existence in binary systems imply that these may possess rapid rotation rates (Bhattacharya & van den Heuvel 1991 and references therein). Particularly, the two SS candidates in SAX J1808.4 –3658 and 4U 1728 –34

are millisecond pulsars having spin periods $P = 2.49$ ms and $P = 2.75$ ms respectively. These two properties make the incorporation of general relativistic effects of rotation imperative for satisfactory treatment of the problem.

Most of the calculations on the rotational properties of SS, reported so far, have relied on the slow rotation approximation (Colpi & Miller 1992, Glendenning & Weber 1992). This approximation loses its validity as the star’s spin frequency approaches the mass shedding limit. Rapidly rotating SS sequences have been recently computed by Gourgoulhon *et al.* (1999), Stergioulas *et al.* (1999), and Zdunik *et al.* (2000). However, all the calculations mentioned above make use of a very schematic model (Freedman & McLerran 1978, Farhi & Jaffe, 1984), related to the MIT bag model (Chodos *et al.* 1974) for hadrons, for the equation of state (EOS) of SQM. Within the MIT bag model EOS, the SS radii calculated are seen to be incompatible with the mass–radius (M – R) relation (Li *et al.* 1999a) for SAX J1808.4 –3658, and only marginally compatible (see Fig. 2) with that for 4U 1728 –34 (Li *et al.* 1999b).

In this letter, we present calculations of equilibrium sequences of rapidly rotating SS in general relativity using a new model for the EOS of SQM derived by Dey *et al.* (1998). This model is based on a “dynamical” density–dependent approach to confinement. In contrast, in the simple EOS based on the MIT bag model, the medium effects on the quark degrees of freedom and on the quark–quark interaction are not considered. For illustrative purposes, we compare our results, obtained using this new EOS, with those for the MIT bag model EOS.

We use the methodology described in detail in Datta, Thampan & Bombaci (1998) to calculate the structure of rapidly rotating SS. For completeness, we briefly describe the method here. For a general axisymmetric and stationary space–time, assuming a perfect fluid configuration, the Einstein field equations reduce to ordinary integrals (using Green’s function approach). These integrals may be self consistently (numerically and iteratively) solved to yield

¹Dipartimento di Fisica, Università di Pisa, and INFN Sezione Pisa, via Buonarroti 2, I–56127, Italy; BOMBACI@pisa.infn.it

²Inter–University Centre for Astronomy and Astrophysics (IUCAA), Pune 411 007, India; arun@iucaa.ernet.in

³Indian Institute of Astrophysics, Bangalore 560034, India

⁴Raman Research Institute, Bangalore 560080, India

the value of metric coefficients in all space. Using these metric coefficients, one may then compute the structure parameters, angular momentum and moment of inertia corresponding to initially assumed central density and polar to equatorial radius ratio. These may then be used (as described in Thampan & Datta 1998) to calculate parameters connected with stable circular orbits (like the innermost stable orbit and the Keplerian angular velocities) around the configuration in question.

The sequences that we calculate are: constant rest mass sequences, constant angular velocity sequences, constant central density sequences and constant angular momentum sequences. We also calculate the radius r_{orb} of the marginally stable orbit and its dependence on the spin rate of the SS, which will be relevant for modeling X-ray burst sources involving SS.

2. THE EQUATION OF STATE FOR SQM

As mentioned earlier, the schematic EOS for SQM based on the MIT bag model, has become the “standard” EOS model for SS studies. However, this EOS model becomes progressively less trustworthy as one goes from very high density region (asymptotic freedom regime) to that of low densities, where confinement (hadron formation) takes place. Recently, Dey *et al.* (1998) derived a new EOS for SQM using a “dynamical” density-dependent approach to confinement. This EOS has asymptotic freedom built in, shows confinement at zero baryon density, deconfinement at high density. In this model, the quark interaction is described by a colour-Debye-screened inter-quark vector potential originating from gluon exchange, and by a density-dependent scalar potential which restores chiral symmetry at high density (in the limit of massless quarks). This density-dependent scalar potential arises from the density dependence of the in-medium effective quark masses M_q , which are taken to depend on the baryon number density n_B according to (see Dey *et al.* 1998)

$$M_q = m_q + 310 \cdot \text{sech}\left(\nu \frac{n_B}{n_0}\right) \quad (\text{MeV}), \quad (1)$$

where $n_0 = 0.16 \text{ fm}^{-3}$ is the normal nuclear matter density, $q (= u, d, s)$ is the flavor index, and ν is a parameter. The effective quark mass $M_q(n_B)$ goes from its constituent mass at $n_B = 0$, to its current mass m_q , as $n_B \rightarrow \infty$. Here, we consider two different parameterizations of this EOS, which correspond to a different choice for the parameter ν . The equation of state SS1 (SS2) corresponds to $\nu = 0.333$ ($\nu = 0.286$). These two models for the EOS give absolutely stable SQM according to the strange matter hypothesis.

In order to compare our results with those of previous studies, we also use the MIT bag model EOS, for massless non-interacting quarks and $B = 90 \text{ MeV/fm}^3$ (hereafter the $B90_0$ EOS).

3. RESULTS AND DISCUSSION

The equilibrium sequences of rotating SS depend on two parameters: the central density (ρ_c) and the rotation rate (Ω). For purpose of illustration, we choose three limits in this parameter space. These are: (i) the static or non-rotating limit, (ii) the limit at which instability to quasi-radial mode sets in and (iii) the centrifugal mass shed limit. The last limit corresponds to the maximum Ω for which

centrifugal forces are able to balance the inward gravitational force.

The result of our calculations for EOS SS1 is displayed in Fig. 1. In Fig. 1 (a) we show the functional dependence of the gravitational mass (M) with ρ_c . In these set of figures, the bold solid curve represents the non-rotating or static limit, and the bold dashed curve the centrifugal mass shed limit. The thin solid curves are the constant rest mass (M_0) evolutionary sequences. The evolutionary sequences above the maximum stable non-rotating mass configuration are the supramassive evolutionary sequences, and those that lie below this limit are the normal evolutionary sequences. The maximum mass sequence for this EOS corresponds to $M_0 = 2.2 M_\odot$. The thin dot-dashed line (slanted towards left) represents instability to quasi-radial perturbations. In the central panel of the same figure, we give a plot of M as a function of the equatorial radius R .

In panel (c) of Fig. 1, we display the plot of Ω as a function of the specific angular momentum $\tilde{j} = cJ/GM_0^2$ (where J is the angular momentum of the configuration). Unlike for neutron stars (e.g. Cook, Shapiro & Teukolsky 1994; Datta, Thampan & Bombaci 1998), the Ω - \tilde{j} curve does not show a turn-over to lower \tilde{j} values for SS. This is due to the effect of the long-range (non-perturbative) interaction in QCD (Quantum Chromo-Dynamics), which is responsible for quark confinement in hadrons, and makes low mass SS self-bound objects. Ω for the mass shed limit appears to asymptotically tend to a non-zero value for rapidly rotating low mass stars. A further ramification of this result is that the ratio of the rotational energy to the total gravitational energy (T/W) becomes greater than 0.21 (as also reported by Gourgoulhon *et al.* 1999) thus probably making the configurations susceptible to triaxial instabilities.

In Table 1 we display the values of the structure parameters for the maximum mass non-rotating SS models. The larger value of the maximum mass for the SS1 model, with respect to the SS2 model, can be traced back to role of the parameter ν in eq. (1) for the effective quark mass M_q . In fact, a larger value of ν (SS1 model) gives a faster decrease of M_q with density, producing a stiffer EOS.

Table 2 and Table 3 displays the maximum mass rotating and maximum angular momentum models for the EOS models under consideration. While for EOS SS1, the maximum mass rotating model and the maximum angular momentum models are the same, for EOS SS2, the two models are slightly different, with the maximum angular momentum model coming earlier (with respect to ρ_c) than the maximum mass rotating configuration.

In Table 4 we list the values of the various parameters for the constant Ω sequences for EOS SS1. The first entry in this table corresponds to ρ_c for which $r_{\text{orb}} = R$. For higher values of ρ_c , $r_{\text{orb}} > R$; for large values of ρ_c , the boundary layer (separation between the surface of the SS and its innermost stable orbit) can be substantial ($\sim 5 \text{ km}$ for the maximum value of the listed ρ_c).

In Fig. 2, we plot our theoretically calculated M - R curves (dashed curves) for EOS SS1 and B90_0, for a rotational frequency of 364 Hz corresponding to the inferred rotational frequency of the compact star in the source 4U 1728-34 (Méndez & Van der Klis 1999). In the same figure, we plot the radius R_0 of the inner edge of the ac-

cretion disk (dotted curve) for 4U 1728 –34 as deduced by Titarchuk & Osherovich (1999) (see also Li et al. 1999b) from a fit of kHz QPO data in this source. Since R_0 must be larger than both R and r_{orb} , it is possible to deduce (Li et al. 1999b) an upper bound for the radius and mass of the central accretor. Using these constraints on M and R , Li et al. (1999b) concluded that 4U 1728 –34 is possibly a strange star rather than a neutron star. However, in their calculation, Li et al. (1999b) used an approximate EOS-independent expression to account for the effects of rotation on the moment of inertia of the star and hence on r_{orb} (see Li et al 1999b for further details). In contrast, in this letter we make an “exact” calculation (as a result of our accurate calculation of angular momentum and hence the moment of inertia) of r_{orb} (triple-dot dashed curves) for the two EOS SS1 and B90_0. Notice that the upper bound for the mass of 4U 1728 –34 (the intersection point between the curves R_0 and r_{orb}) is about $1.15 M_{\odot}$, in agreement with the results of Li et al. (1999b). The two EOS models considered in Fig. 2 are both consistent with

4U 1728 –34. Also, having in mind the scaling with $B^{-1/2}$ of the M–R relation of SS within the bag model EOS (Witten, 1984) and the constraints on the allowed values of the constant B to fulfil the strange matter hypothesis (Farhi & Jaffe, 1984), we see that the simple EOS based on the MIT bag model is only marginally compatible with the M–R relation for 4U 1728 –34.

To summarize, in this letter, we present calculations of equilibrium sequences of rapidly rotating strange stars in general relativity, for a set of new equations of state. We compare the results so obtained with those for the generally used MIT bag model EOS. In addition to this, we illustrate the results of our computations for a specific rotation rate inferred for the central accretor in the source 4U 1728 –34. We find that if the compact star in this source were indeed to be a strange star, then its mass and radius would be bounded above by a value of $M = 1.15 M_{\odot}$ and $R = 9.3$ km. It is expected that future observations may shed more light on this issue.

REFERENCES

- Alcock, C., Farhi, E., & Olinto, A. 1986, ApJ, 310, 261
 Backer, D.C. *et al.* 1982, Nature, 300, 615
 Bhattacharya, D., & van den Heuvel, E.P.J. 1991, Phys. Rep. 203, 1
 Bodmer, A.R. 1971, Phys. Rev. D, 4, 1601
 Bombaci, I. 1997, Phys. Rev. C, 55, 1587
 Cheng, K.S., Dai, Z.G., Wai, D.M., & Lu, T., 1998, Science, 280, 407
 Chodos, A., Jaffe, R.L., Johnson, K., Thorn, C.B., & Weisskopf, V.F. 1974, Phys. Rev. D9, 3471
 Colpi, M. & Miller, J.C. 1992, ApJ, 388 513
 Cook, G.B., Shapiro, S.L. & Teukolsky, S.A. 1994, ApJ, 424,823
 Datta, B., Thampan, A.V., & Bombaci, I. 1998, A&A, 334, 943
 Dey, M., Bombaci, I., Dey, J., Ray, S., & Samanta, B.C. 1998, Phys. Lett. B, 438, 123; erratum 1999, Phys. Lett. B, 467, 303
 Farhi, E., & Jaffe, R.L. 1984, Phys. Rev. D30, 2379
 Freedman, B. & McLerran, L. 1978, Phys. Rev. D17, 1109
 Glendenning, N.K. & Weber F. 1992, ApJ, 400, 647
 Gourgoulhon, E. *et al.* 1999, A&A, 349, 851
 Haensel, P., Zdunik, J.L. & Schaefer, R. 1986, A&A, 160, 121
 Li, X.-D., Bombaci, I., Dey, M., Dey J., & van den Heuvel, E.P.J. 1999a, Phys. Rev. Lett., 83, 3776
 Li, X.-D., Ray, S., Dey, J., Dey, M., & Bombaci, I. 1999b, ApJ, 527, L51
 Méndez, M. & Van der Klis, M. 1999, ApJ, 517, L51
 Stergioulas, N., Kluźniak, W. & Bulik, T. 1999, A&A, 352, L116
 Terazawa, H. 1979, INS-Report, 336 (INS, Univ. of Tokyo); 1989, J. Phys. Soc. Japan, 58, 3555; 1989, J. Phys. Soc. Japan, 58, 4388
 Thampan, A.V. & Datta, B. 1998, MNRAS, 297, 570
 Titarchuk, L. & Osherovich, V. 1999, ApJ, 518, L95
 Witten, E. 1984, Phys. Rev. D, 30, 272
 Zdunik, J.L., Haensel, P., Gondek-Rosinska, D. & Gourgoulhon, E. 2000, A&A, 356, 612

TABLE 1

STRUCTURE PARAMETERS FOR THE NON-ROTATING MAXIMUM MASS CONFIGURATIONS. LISTED, ARE THE CENTRAL DENSITY (ρ_c) IN UNITS OF G CM^{-3} , THE GRAVITATIONAL MASS (M) IN SOLAR UNITS, THE EQUATORIAL RADIUS (R) IN KM, THE BARYONIC MASS (M_0) IN SOLAR UNITS, THE RADIUS OF THE MARGINALLY STABLE ORBIT (r_{orb}) IN KM AND THE MOMENT OF INERTIA (I) IN UNITS OF 10^{45} G CM^2 .

EOS	ρ_c	M	R	M_0	r_{orb}	I
SS1	4.65E+15	1.438	7.093	1.880	12.729	0.736
SS2	5.60E+15	1.324	6.533	1.658	11.716	0.578
B90_0	3.09E+15	1.603	8.745	1.937	14.020	1.099

TABLE 2

STRUCTURE PARAMETERS FOR THE MAXIMALLY ROTATING ($\Omega = \Omega_{\text{ms}}$) MAXIMUM MASS CONFIGURATION. IN ADDITION TO THE QUANTITIES LISTED IN THE PREVIOUS TABLE, WE DISPLAY, THE ROTATION RATE (Ω) IN 10^4 RAD S^{-1} , THE RATIO OF THE ROTATIONAL TO THE TOTAL GRAVITATIONAL ENERGY (T/W) AND THE SPECIFIC ANGULAR MOMENTUM ($\tilde{j} = cJ/GM_0^2$).

EOS	ρ_c	Ω	I	M	T/W	R	\tilde{j}	r_{orb}	M_0
SS1	3.10E+15	1.613	2.072	2.077	0.219	10.404	0.524	11.656	2.694
SS2	3.60E+15	1.738	1.613	1.904	0.218	9.612	0.570	10.758	2.366
B90_0	1.90E+15	1.190	3.369	2.272	0.232	13.213	0.633	14.612	2.683

TABLE 3

STRUCTURE PARAMETERS FOR THE MAXIMUM ANGULAR MOMENTUM CONFIGURATION.

EOS	ρ_c	Ω	I	M	T/W	R	\tilde{j}	r_{orb}	M_0
SS1	3.10E+15	1.613	2.072	2.077	0.219	10.404	0.524	11.656	2.694
SS2	3.40E+15	1.719	1.633	1.899	0.220	9.693	0.575	10.837	2.355
B90_0	1.70E+15	1.161	3.456	2.254	0.239	13.447	0.650	14.864	2.650

TABLE 4

STRUCTURE PARAMETERS FOR THE CONSTANT ANGULAR VELOCITY SEQUENCE FOR EOS SS1. THIS SEQUENCE CORRESPONDS TO THE ROTATION RATE OF THE PULSAR PSR 1937 +21 (BACKER *et al.* 1982), HAVING $\Omega = 4.03 \times 10^3 \text{ RAD S}^{-1}$ OR PERIOD $P = 1.556 \text{ MS}$.

ρ_c	I	M	T/W	R	\tilde{j}	r_{orb}	M_0
1.70E+15	0.353	0.852	0.013	6.663	0.153	6.663	1.027
1.80E+15	0.433	0.963	0.012	6.884	0.143	7.664	1.178
1.90E+15	0.502	1.052	0.019	7.036	0.136	8.378	1.301
2.40E+15	0.701	1.297	0.010	7.326	0.116	10.355	1.660
2.60E+15	0.737	1.346	0.010	7.344	0.112	10.741	1.735
4.60E+15	0.769	1.458	0.008	7.139	0.096	11.732	1.914
5.65E+15	0.734	1.449	0.007	7.007	0.093	11.703	1.899

FIG. 1.— Structure parameters for rotating strange stars corresponding to EOS SS1. The bold-solid line represents the non-rotating limit, the bold-dotted line the mass-shed limit and the almost vertical thin dot-dashed line tilted to the left is the instability limit to quasi-radial mode perturbations. The thin solid lines (labelled 1, 2 ...) represent constant rest mass sequences: 1: $1.59 M_{\odot}$, 2: $1.66 M_{\odot}$, 3: $1.88 M_{\odot}$, 4: $2.14 M_{\odot}$ 5: $2.41 M_{\odot}$.




FIG. 2.— Mass–radius relations and r_{orb} for a rotation rate of $\Omega = 2.287 \times 10^3 \text{ rad s}^{-1}$ (i.e. 364 Hz) corresponding to the inferred rotation rate of the central accretor in 4U 1728 –34 (Médéz and Van der Klis 1999). Also plotted are the bounds on the M – R from general considerations, and the fit for these values obtained by Titarchuk & Osherovich (1999). See text for further details.

